

AD-A225 250

## REPORT DOCUMENTATION PAGE

Form Approved

OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 12 Aug 90		3. REPORT TYPE AND DATES COVERED INTERIM - April 1989 - September 1989	
4. TITLE AND SUBTITLE ULTRASTRUCTURAL STUDY OF THE EFFECT OF POLYCTFE ON PRIMATE LIVER				5. FUNDING NUMBERS PE 62202F PR 6302 TA 630201 WU 63020169  Accession No. 316065	
6. AUTHOR(S) Mattie, D.R., Maslanka, J.J., and Chase, M.R.				8. PERFORMING ORGANIZATION REPORT NUMBER  AAMRL-TR-90-037	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) AAMRL, Toxic Hazards Division HSD, AFSC Wright-Patterson AFB, Ohio 45433-6573				9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)  AUG 16 1990	
11. SUPPLEMENTARY NOTES Proceedings of the XIIth International Congress for Electron Microscopy, pp 280-281, 1990				10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
12a. DISTRIBUTION/AVAILABILITY STATEMENT  Approved for Public Release; distribution is Unlimited				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Oligomers of chlorotrifluoroethylene (POLYCTFE) have been found to possess excellent properties as lubricants and hydraulic fluids. The 3.1 oil, consisting primarily of oligomers of 3 and 4 units, is a candidate base stock for nonflammable hydraulic fluids for advanced aircraft. Unfortunately, ultrastructural studies of the livers from rats exposed to vapors of POLYCTFE for 90-days revealed an increase in the number of peroxisomes, dose-dependent mitochondrial swelling and a dose-dependent increase in smooth endoplasmic reticulum. In order to determine the relevance of the effect of POLYCTFE on rat livers to human livers, four male rhesus monkeys weighing 8 to 10 kg were dosed by gavage daily for 15 days with 0.725 g/kg POLYCTFE. Wedge biopsies of liver obtained from each animal were collected and prepared for transmission electron microscopic examination. Repeated exposure of primates to POLYCTFE for 15-days resulted in only minor changes in the hepatocytes. There was no evidence of peroxisomal proliferation in exposed primate livers. Oligomers of CTFE did not produce the same effect in primate livers as in rat livers. The toxicity of POLYCTFE in rats may not be an appropriate indicator of potential human hazard.					
14. SUBJECT TERMS (U) Toxicity (U) Primates (U) Liver (U) Halogenated Hydrocarbons (U) Electron Microscopy, 325				15. NUMBER OF PAGES 2	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED				16. PRICE CODE	
18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED		19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED		20. LIMITATION OF ABSTRACT UL	

The Report Documentation Page (RDP) is used in announcing and cataloging reports. It is important that this information be consistent with the rest of the report, particularly the cover and title page. Instructions for filling in each block of the form follow. It is important to *stay within the lines* to meet optical scanning requirements.

**Block 1. Agency Use Only (Leave blank).**

**Block 2. Report Date.** Full publication date including day, month, and year, if available (e.g. 1 Jan 88). Must cite at least the year.

**Block 3. Type of Report and Dates Covered.** State whether report is interim, final, etc. If applicable, enter inclusive report dates (e.g. 10 Jun 87 - 30 Jun 88).

**Block 4. Title and Subtitle.** A title is taken from the part of the report that provides the most meaningful and complete information. When a report is prepared in more than one volume, repeat the primary title, add volume number, and include subtitle for the specific volume. On classified documents enter the title classification in parentheses.

**Block 5. Funding Numbers.** To include contract and grant numbers; may include program element number(s), project number(s), task number(s), and work unit number(s). Use the following labels:

C - Contract	PR - Project
G - Grant	TA - Task
PE - Program Element	WU - Work Unit Accession No.

**Block 6. Author(s).** Name(s) of person(s) responsible for writing the report, performing the research, or credited with the content of the report. If editor or compiler, this should follow the name(s).

**Block 7. Performing Organization Name(s) and Address(es).** Self-explanatory.

**Block 8. Performing Organization Report Number.** Enter the unique alphanumeric report number(s) assigned by the organization performing the report.

**Block 9. Sponsoring/Monitoring Agency Name(s) and Address(es).** Self-explanatory.

**Block 10. Sponsoring/Monitoring Agency Report Number.** (If known)

**Block 11. Supplementary Notes.** Enter information not included elsewhere such as: Prepared in cooperation with...; Trans. of...; To be published in.... When a report is revised, include a statement whether the new report supersedes or supplements the older report.

**Block 12a. Distribution/Availability Statement.** Denotes public availability or limitations. Cite any availability to the public. Enter additional limitations or special markings in all capitals (e.g. NOFORN, REL, ITAR).

DOD - See DoDD 5230.24, "Distribution Statements on Technical Documents."

DOE - See authorities.

NASA - See Handbook NHB 2200.2.

NTIS - Leave blank.

**Block 12b. Distribution Code.**

DOD - Leave blank.

DOE - Enter DOE distribution categories from the Standard Distribution for Unclassified Scientific and Technical Reports.

NASA - Leave blank.

NTIS - Leave blank.

**Block 13. Abstract.** Include a brief (Maximum 200 words) factual summary of the most significant information contained in the report.

**Block 14. Subject Terms.** Keywords or phrases identifying major subjects in the report.

**Block 15. Number of Pages.** Enter the total number of pages.

**Block 16. Price Code.** Enter appropriate price code (NTIS only).

**Blocks 17. - 19. Security Classifications.** Self-explanatory. Enter U.S. Security Classification in accordance with U.S. Security Regulations (i.e., UNCLASSIFIED). If form contains classified information, stamp classification on the top and bottom of the page.

**Block 20. Limitation of Abstract.** This block must be completed to assign a limitation to the abstract. Enter either UL (unlimited) or SAR (same as report). An entry in this block is necessary if the abstract is to be limited. If blank, the abstract is assumed to be unlimited.

## ULTRASTRUCTURAL STUDY OF THE EFFECT OF POLYCTFE ON PRIMATE LIVER

Mattie, D.R., Maslanka, J.J., and Chase, M.R.

Toxic Hazards Division, Harry G. Armstrong Aerospace Medical Research Laboratory  
(AAMRL/TH), WPAFB, OH 45433-6573

Polychlorotrifluoroethylene (POLYCTFE) existing as oligomers of 3 and 4 units, is a candidate base stock for a nonflammable hydraulic fluid for advanced aircraft. POLYCTFE caused significant changes in the livers of rats after repeated inhalation exposure.<sup>1</sup> Ultrastructural studies of the livers from rats exposed to vapors of POLYCTFE oligomers for 90-days revealed an increase in the number of peroxisomes, dose-dependent mitochondrial swelling and a dose-dependent increase in smooth endoplasmic reticulum.<sup>2</sup> The objective of this study was to ultrastructurally examine hepatocytes from the livers of primates exposed to POLYCTFE and compare the exposed primate hepatocytes to those from rat livers, in order to determine the relevance of the effect of POLYCTFE on rat livers to human livers.

Male rhesus monkeys weighing 8 to 10 kg were dosed by gavage daily for 15 days with 0.725 g/kg POLYCTFE. Four animals received POLYCTFE and 3 received a comparable amount of water. A laparotomy was performed on each animal at the end of the dosing period in order to obtain a wedge biopsy for analysis. A 1mm slice of the liver biopsy from each animal was collected for transmission electron microscopic examination. The liver slices were fixed in 2% glutaraldehyde in 0.1M cacodylate buffer at pH 7.4 and minced into 1 cubic mm pieces. The minced tissue was post-fixed with 2% osmium and was processed into EPOX 812 plastic capsules. One micron thick sections were cut in order to identify centrilobular zones. Thin sections were cut from the centrilobular and intermediate zones of liver lobules. Thin sections stained with uranyl acetate and lead citrate, were examined with a JEOL 100B electron microscope at 60 kV. For each animal 20-30 photographs were taken of 3 or more hepatocytes that were representative of the liver for that animal. The number of peroxisomes were counted per 30,000 magnification visual field using an 8 by 10 inch photograph. An average of 9 photographs were counted per each animal. Statistical analysis of peroxisome counts was conducted using the Student's T-Test at the  $p < 0.05$  level of significance.

Repeated exposure of primates to POLYCTFE for 15-days resulted in only minor changes in the hepatocytes with no evidence of peroxisomal proliferation (FIG. 1,2). The mean number of peroxisomes for control primate liver was  $2.5 \pm 0.33$  ( $\pm$  S.E.M.) compared to  $3.3 \pm 0.42$  for exposed animals. The difference between the control and exposed values for peroxisomal counts were not statistically significant. Mild to moderate mitochondrial swelling was seen in both control (FIG. 3) and exposed animals (FIG. 4). A decrease in glycogen was seen in primates exposed to POLYCTFE (FIG. 2 or 4). POLYCTFE did not produce the same effect in primate livers as in rat livers.<sup>2</sup> Mitochondrial swelling seen in both control and exposed animals was probably a result of ischemia due to the method of obtaining the biopsy.<sup>3</sup> An observed decrease in food consumption by all animals could explain the decreased amount of glycogen in exposed animals. POLYCTFE did not produce the same effect in primate livers as seen in rat livers.

1. E.R. Kinkead et al., The Toxicologist 9(1989)144.
2. D.R. Mattie et al., Proc. Ann. EMSA Meeting 46(1988)330.
3. Dougherty, C., Personal Communication, 1989.



FIG. 1.--Hepatocyte from control primate liver cell showing normal number of peroxisomes (P). Bar = 1  $\mu$ m.

FIG. 2.--Hepatocyte from primate liver exposed to CTFE for 15-days showing normal number of peroxisomes (P). Bar = 1  $\mu$ m.

FIG. 3.--Control primate hepatocytes with glycogen (G) and swollen mitochondria (M). Bar = 1  $\mu$ m.

FIG. 4.--Primate hepatocyte from liver exposed to CTFE for 15-days with swollen mitochondria (M) and showing a decreased amount of glycogen (G). Bar = 1  $\mu$ m.



Dist	Special	
A-1	20	